



# 6 FUNCTIONS



## Let's Study

- Function, Domain, Co-domain, Range
- Types of functions
- Representation of function
- Basic types of functions
- Piece-wise defined and special functions



## Let's : Learn

### 6.1 Function

**Definition :** A function (or mapping)  $f$  from a set  $A$  to set  $B$  ( $f: A \rightarrow B$ ) is a relation which associates for each element  $x$  in  $A$ , a unique (exactly one) element  $y$  in  $B$ .

Then the element  $y$  is expressed as  $y = f(x)$ .

$y$  is the image of  $x$  under  $f$ .

$f$  is also called a map or transformation.

If such a function exists, then  $A$  is called the **domain** of  $f$  and  $B$  is called the **co-domain** of  $f$ .

### Illustration:

Examine the following relations which are given by arrows of line segments joining elements in  $A$  and elements in  $B$ .




Fig. 6.1

Since, every element from  $A$  is associated to exactly one element in  $B$ ,  $R_1$  is a well defined function.




Fig. 6.2

$R_2$  is not a function because element 'd' in  $A$  is not associated to any element in  $B$ .




Fig. 6.3

$R_3$  is not a function because element  $a$  in  $A$  is associated to two elements in  $B$ .

The relation which defines a function  $f$  from domain  $A$  to co-domain  $B$  is often given by an algebraic rule.

For example,  $A = Z$ , the set of integers and  $B = Q$  the set of rational numbers and the function

$f$  is given by  $f(n) = \frac{n}{7}$  here  $n \in Z, f(n) \in Q$ .

### 6.1.1 Types of function

#### One-one or One to one or Injective function

**Definition :** A function  $f : A \rightarrow B$  is said to be one-one if different elements in A have different images in B. The condition is also expressed as

$$f(a) = f(b) \Rightarrow a = b \quad [\text{As } a \neq b \Rightarrow f(a) \neq f(b)]$$

#### Onto or Surjective function

**Definition:** A function  $f : A \rightarrow B$  is said to be onto if every element  $y$  in B is an image of some  $x$  in A (or  $y$  in B has preimage  $x$  in A)

The image of A can be denoted by  $f(A)$ .


$$f(A) = \{y \in B \mid y = f(x) \text{ for some } x \in A\}$$

$f(A)$  is also called the **range** of  $f$ .

Note that  $f : A \rightarrow B$  is onto if  $f(A) = B$ .


Also range of  $f = f(A) \subset$  co-domain of  $f$ .

#### Illustration:




**Fig. 6.4**

$f_1$  is one-one, but not onto as element 5 is in B has no pre image in A




**Fig. 6.5**

$f_2$  is one-one, and onto



**Fig. 6.6**

$f_3$  is onto but not one-one as  $f(a) = f(b) = 1$  but  $a \neq b$ .





**Fig. 6.7**

$f_4$  is neither one-one, nor onto

### 6.1.2 Representation of Function


Verbal form	Output exceeds twice the input by 1 Domain : Set of inputs Range : Set of outputs
Arrow form on Venn Diagram	<p><b>Fig. 6.8</b></p>
Ordered Pair (x, y)	Domain : Set of pre-images Range: Set of images $f = \{(2,5), (3,7), (4,9), (5,11)\}$ Domain : Set of 1 <sup>st</sup> components from each ordered pair = $\{2, 3, 4, 5\}$ Range : Set of 2 <sup>nd</sup> components from each ordered pair = $\{5, 7, 9, 11\}$

Rule / Formula	$y = f(x) = 2x + 1$ Where $x \in N, 1 < x < 6$ $f(x)$ read as 'f of x' or 'function of x' Domain : Set of values of $x$ for which $f(x)$ is defined Range : Set of values of $y$ for which $f(x)$ is defined										
Tabular Form	<table border="1" style="display: inline-table; margin-right: 20px;"> <thead> <tr><th><math>x</math></th><th><math>y</math></th></tr> </thead> <tbody> <tr><td>2</td><td>5</td></tr> <tr><td>3</td><td>7</td></tr> <tr><td>4</td><td>9</td></tr> <tr><td>5</td><td>11</td></tr> </tbody> </table> Domain : $x$ values Range: $y$ values	$x$	$y$	2	5	3	7	4	9	5	11
$x$	$y$										
2	5										
3	7										
4	9										
5	11										
Graphical form	 <p style="text-align: center;"><b>Fig. 6.9</b></p> Domain: Projection of graph on $x$ -axis. Range: Projection of graph on $y$ -axis.										



**Fig. 6.10**

Since every  $x$  has a unique associated value of  $y$ .  
 It is a function.




**Fig. 6.11**

This graph does not represent a function as vertical line intersects at more than one point some  $x$  has more than one values of  $y$ .

**Horizontal Line Test:**

If no horizontal line intersects the graph of a function in more than one point, then the function is one-one function.

**Illustration:**



**Fig. 6.12**

**6.1.3 Graph of a function:**


If the domain of function is in  $R$ , we can show the function by a graph in  $xy$  plane. The graph consists of points  $(x,y)$ , where  $y = f(x)$ .

**Vertical Line Test**

Given a graph, let us find if the graph represents a function of  $x$  i.e.  $f(x)$

A graph represents function of  $x$ , only if no vertical line intersects the curve in more than one point.

The graph is a one-one function as a horizontal line intersects the graph at only one point.



**Fig. 6.13**

The graph is a one-one function

**6.1.4 Value of function :**  $f(a)$  is called the value of function  $f(x)$  at  $x = a$

**Evaluation of function:**


**Ex. 1)** Evaluate  $f(x) = 2x^2 - 3x + 4$  at  $x = 7$  &  $x = -2t$

**Solution :**  $f(x)$  at  $x = 7$  is  $f(7)$

$$\begin{aligned} f(7) &= 2(7)^2 - 3(7) + 4 \\ &= 2(49) - 21 + 4 \\ &= 98 - 21 + 4 \\ &= 81 \end{aligned}$$

$$\begin{aligned} f(-2t) &= 2(-2t)^2 - 3(-2t) + 4 \\ &= 2(4t^2) + 6t + 4 \\ &= 8t^2 + 6t + 4 \end{aligned}$$

**Ex. 2)** Using the graph of  $y = g(x)$ , find  $g(-4)$  and  $g(3)$



**Fig. 6.14**

**Solution :** From graph when  $x = -4$ ,  $y = 0$   
so  $g(-4) = 0$

From graph when  $x = 3$ ,  $y = -5$  so  $g(3) = -5$

**Function Solution:**


**Ex. 3)** If  $t(m) = 3m^2 - m$  and  $t(m) = 4$ , then find  $m$

**Solution :** As

$$\begin{aligned} t(m) &= 4 \\ 3m^2 - m &= 4 \\ 3m^2 - m - 4 &= 0 \\ 3m^2 - 4m + 3m - 4 &= 0 \\ m(3m - 4) + 1(3m - 4) &= 0 \\ (3m - 4)(m + 1) &= 0 \end{aligned}$$

Therefore,  $m = \frac{4}{3}$  or  $m = -1$


**Ex. 4)** From the graph below find  $x$  for which  $f(x) = 4$



**Fig. 6.15**

**Solution :** To solve  $f(x) = 4$  i.e.  $y = 4$

Find the values of  $x$  where graph intersects line  $y = 4$



**Fig. 6.16**

Therefore,  $x = -1$  and  $x = 3$ .

**Function from equation:**

**Ex. 5 (Activity)** From the equation  $4x + 7y = 1$  express

- i)  $y$  as a function of  $x$
- ii)  $x$  as a function of  $y$

**Solution :** Given equation is  $4x + 7y = 1$

- i) From the given equation

$$7y = \square$$

$$y = \square = \text{function of } x$$

$$\text{So } y = f(x) = \square$$

- ii) From the given equation

$$4x = \square$$

$$x = \square = \text{function of } y$$

$$\text{So } x = g(y) = \square$$


**6.1.5 Some Basic Functions**

(Here  $f: \mathbb{R} \rightarrow \mathbb{R}$  Unless stated otherwise)

**1) Constant Function**

**Form :**  $f(x) = k, k \in \mathbb{R}$

**Example :** Graph of  $f(x) = 2$




**Fig. 6.17**

**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $\{2\}$

**2) Identity function**

If  $f: \mathbb{R} \rightarrow \mathbb{R}$  then identity function is defined by  $f(x) = x$ , for every  $x \in \mathbb{R}$ .

Identity function is given in the graph below.



**Fig. 6.18**

**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $\mathbb{R}$  or  $(-\infty, \infty)$


[**Note :** Identity function is also given by  $I(x) = x$ ].

**3) Power Functions :  $f(x) = ax^n, n \in \mathbb{N}$**

(Note that this function is a multiple of  $n^{\text{th}}$  power of  $x$ )

**i) Square Function**

**Example :**  $f(x) = x^2$



**Fig. 6.19**


**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $[0, \infty)$

**Properties:**

- 1) Graph of  $f(x) = x^2$  is a parabola opening upwards and with vertex at origin.
- 2) Graph is symmetric about  $y$  - axis .
- 3) The graph of even powers of  $x$  looks similar to square function. (verify !) e.g.  $x^4, x^6$ .
- 4)  $(y - k) = (x - h)^2$  represents parabola with vertex at  $(h, k)$
- 5) If  $-2 \leq x \leq 2$  then  $0 \leq x^2 \leq 4$  (see fig.) and if  $-3 \leq x \leq 2$  then  $0 \leq x^2 \leq 9$  (see fig).

**ii) Cube Function**

**Example :**  $f(x) = x^3$



**Fig. 6.20**

**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $\mathbb{R}$  or  $(-\infty, \infty)$

**Properties:**

- 1) The graph of odd powers of  $x$  (more than 1) looks similar to cube function. e.g.  $x^5, x^7$ .

**4) Polynomial Function**


$$f(x) = a_0 x^n + a_1 x^{n-1} + \dots + a_{n-1} x + a_n$$

is polynomial function of degree  $n$  , if  $a_0 \neq 0$ , and  $a_i$  s are real.

**i) Linear Function**

**Form :**  $f(x) = ax + b$  ( $a \neq 0$ )

**Example :**  $f(x) = -2x + 3, x \in \mathbb{R}$



**Fig. 6.21**


**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $\mathbb{R}$  or  $(-\infty, \infty)$

**Properties :**

- 1) Graph of  $f(x) = ax + b$  is a line with slope 'a',  $y$ -intercept 'b' and  $x$ -intercept  $\left(-\frac{b}{a}\right)$ .
- 2) Function : is increasing when slope is positive and decreasing when slope is negative.

**ii) Quadratic Function**

**Form :**  $f(x) = ax^2 + bx + c$  ( $a \neq 0$ )




**Fig. 6.22**

**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $[k, \infty)$

**Properties :**

- 1) Graph of  $f(x) = ax^2 + bx + c$  and where  $a > 0$  is a parabola.



**Fig. 6.23**

Consider,  $y = ax^2 + bx + c$

$$= a \left( x^2 + \frac{b}{a}x + \frac{b^2}{4a^2} \right) + c - \frac{b^2}{4a}$$

$$= a \left( x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a}$$

$$\left( y + \frac{b^2 - 4ac}{4a} \right) = a \left( x + \frac{b}{2a} \right)^2$$

With change of variable

$$X = x + \frac{b}{2a}, Y = y + \frac{b^2 - 4ac}{4a}$$

this is a parabola  $Y = aX^2$

This is a parabola with vertex

$$\left( -\frac{b}{2a}, \frac{b^2 - 4ac}{4a} \right) \text{ or } \left( \frac{-b}{2a}, \frac{-D}{4a} \right) \text{ where}$$

$D = b^2 - 4ac$  and the parabola is opening upwards.

There are three possibilities.

For  $a > 0$ ,

- i) If  $D = b^2 - 4ac = 0$ , the parabola touches x-axis and  $y \geq 0$  for all  $x$ . e.g.  $g(x) = x^2 - 2x + 1$
- ii) If  $D = b^2 - 4ac > 0$ , then parabola intersects x-axis at 2 distinct points. Here  $y$  is negative for values of  $x$  between the 2 roots and positive for large or small  $x$ .


- iii) If  $D = b^2 - 4ac < 0$ , the parabola lies above x-axis and  $y \neq 0$  for any  $x$ . Here  $y$  is positive for all values of  $x$ . e.g.  $f(x) = x^2 + 4x + 5$

### iii) Cubic Function

**Example :**  $f(x) = ax^3 + bx^2 + cx + d$  ( $a \neq 0$ )

**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and

**Range :**  $\mathbb{R}$  or  $(-\infty, \infty)$



**Fig. 6.24**

### Property:

- 1) Graph of  $f(x) = x^3 - 1$

$f(x) = (x - 1)(x^2 + x + 1)$  cuts x-axis at only one point (1,0), which means  $f(x)$  has one real root & two complex roots.

Note that, any polynomial of odd degree must have at least one real root, because the complex roots appear in conjugate pairs.


### 5) Radical Function

**Ex:**  $f(x) = \sqrt[n]{x}$ ,  $n \in \mathbb{N}$

#### 1. Square root function

$$f(x) = \sqrt{x}, x \geq 0$$

(Since square root of negative number is not a real number, so the domain of  $\sqrt{x}$  is restricted to positive values of  $x$ ).



**Fig. 6.25**

**Domain :**  $[0, \infty)$  and **Range :**  $[0, \infty)$

**Note :**

- 1) If  $x$  is positive, there are two square roots of  $x$ . By convention  $\sqrt{x}$  is positive root and  $-\sqrt{x}$  is the negative root.
- 2) If  $-4 < x < 9$ , as  $\sqrt{x}$  is only defined for  $x \geq 0$ , so  $0 \leq \sqrt{x} < 3$ .

**Ex. 6 :** Find the domain and range of  $f(x) = \sqrt{9-x^2}$ .

**Soln. :**  $f(x) = \sqrt{9-x^2}$  is defined for  $9-x^2 \geq 0$ , i.e.  $x^2-9 \leq 0$  i.e.  $(x-3)(x+3) \leq 0$

Therefore  $[-3, 3]$  is domain of  $f(x)$ . (Verify !)

To find range, let  $\sqrt{9-x^2} = y$

Since square root is always positive, so  $y \geq 0$  ... (I)

Also, on squaring we get  $9-x^2 = y^2$

Since,  $-3 \leq x \leq 3$

i.e.  $0 \leq x^2 \leq 9$

i.e.  $0 \geq -x^2 \geq -9$

i.e.  $9 \geq 9-x^2 \geq 9-9$

i.e.  $9 \geq 9-x^2 \geq 0$


i.e.  $3 \geq \sqrt{9-x^2} \geq 0$

$\therefore 3 \geq y \geq 0$  ... (II)

From (I) and (II),  $y \in [0,3]$  is range of  $f(x)$ .

## 2. Cube root function

$$f(x) = \sqrt[3]{x},$$



**Fig. 6.26**

**Domain :**  $\mathbb{R}$  and **Range :**  $\mathbb{R}$

**Note :** If  $-8 \leq x \leq 1$  then  $-2 \leq \sqrt[3]{x} \leq 1$ .

**Ex. 7 :** Find the domain  $f(x) = \sqrt{x^3-8}$ .

**Soln. :**  $f(x)$  is defined for  $x^3-8 \geq 0$

i.e.  $x^3-2^3 \geq 0$ ,  $(x-2)(x^2+2x+4) \geq 0$

In  $x^2+2x+4$ ,  $a=1 > 0$  and  $D=b^2-4ac = 2^2-4 \times 1 \times 4 = -12 < 0$

Therefore,  $x^2+2x+4$  is a positive quadratic.

i.e.  $x^2+2x+4 > 0$  for all  $x$

Therefore  $x-2 \geq 0$ ,  $x \geq 2$  is the domain.

i.e. Domain is  $x \in [2, \infty)$




## 6) Rational Function

**Definition:** Given polynomials

$p(x), q(x)$   $f(x) = \frac{p(x)}{q(x)}$  is defined for  $x$  if  $q(x) \neq 0$ .

**Example :**  $f(x) = \frac{1}{x}, x \neq 0$



**Fig. 6.27**

**Domain :**  $\mathbb{R} - \{0\}$  and **Range :**  $\mathbb{R} - \{0\}$

**Properties:**

- 1) As  $x \rightarrow 0$  i.e. (As  $x$  approaches 0)  $f(x) \rightarrow \infty$  or  $f(x) \rightarrow -\infty$ , so the line  $x = 0$  i.e. y-axis is called vertical asymptote. (A straight line which does not intersect the curve but as  $x$  approaches to  $\infty$  or  $-\infty$  the distance between the line and the curve tends to 0, is called an asymptote of the curve.)
- 2) As  $x \rightarrow \infty$  or  $x \rightarrow -\infty, f(x) \rightarrow 0, y = 0$  the line i.e. x-axis is called horizontal asymptote.
- 3) The domain of rational function  $f(x) = \frac{p(x)}{q(x)}$  is all the real values of  $x$  except the zeroes of  $q(x)$ .

**Ex. 8 :** Find domain and range of the function

$$f(x) = \frac{6 - 4x^2}{4x + 5}$$

**Solution :**  $f(x)$  is defined for all  $x \in \mathbb{R}$  except when denominator is 0.

$$\text{Since, } 4x + 5 = 0 \Rightarrow x = -\frac{5}{4}$$

So Domain of  $f(x)$  is  $\mathbb{R} - \left\{-\frac{5}{4}\right\}$ .

$$\text{To find the range, let } y = \frac{6 - 4x^2}{4x + 5}$$

$$\text{i.e. } y(4x + 5) = 6 - 4x^2$$

$$\text{i.e. } 4x^2 + (4y)x + 5y - 6 = 0.$$

This is a quadratic equation in  $x$  with  $y$  as constant.

Since  $x \in \mathbb{R} - \{-5/4\}$ , i.e.  $x$  is real, we get

Solution if,  $D = b^2 - 4ac \geq 0$

$$\text{i.e. } (4y)^2 - 4(4)(5y - 6) \geq 0$$

$$16y^2 - 16(5y - 6) \geq 0$$

$$y^2 - 5y + 6 \geq 0$$

$$(y - 2)(y - 3) \geq 0$$

Therefore  $y \leq 2$  or  $y \geq 3$  (Verify!)


Range of  $f(x)$  is  $(-\infty, 2] \cup [3, \infty)$

## 7) Exponential Function

**Form :**  $f(x) = a^x$  is an exponential function with base  $a$  and exponent (or index)  $x, a \neq 0,$

$a > 0$  and  $x \in \mathbb{R}.$

**Example :**  $f(x) = 2^x$  and  $f(x) = 2^{-x}$




**Fig. 6.28**

**Domain:**  $\mathbb{R}$  and **Range :**  $(0, \infty)$

**Properties:**

- As  $x \rightarrow -\infty$ , then  $f(x) = 2^x \rightarrow 0$ , so the graph has horizontal asymptote ( $y = 0$ )
- By taking the natural base  $e (\approx 2.718)$ , graph of  $f(x) = e^x$  is similar to that of  $2^x$  in appearance



**Fig. 6.29**

- For  $a > 0, a \neq 1$ , if  $a^x = a^y$  then  $x = y$ . So  $a^x$  is one-one function. (check graph for horizontal line test).
- $r > 1, m > n \Rightarrow r^m > r^n$  and  $r < 1, m > n \Rightarrow r^m < r^n$

**Ex. 9 :** Solve  $5^{2x+7} = 125$ .

**Solution :** As  $5^{2x+7} = 125$

i.e :  $5^{2x+7} = 5^3, \therefore 2x + 7 = 3$

and  $x = \frac{3-7}{2} = \frac{-4}{2} = -2$

**Ex. 10 :** Find the domain of  $f(x) = \sqrt{6 - 2^x - 2^{3-x}}$

**Solution :** Since  $\sqrt{x}$  is defined for  $x \geq 0$

$f(x)$  is defined for  $6 - 2^x - 2^{3-x} \geq 0$

i.e.  $6 - 2^x - \frac{2^3}{2^x} \geq 0$

i.e.  $6 \cdot 2^x - (2^x)^2 - 8 \geq 0$

i.e.  $(2^x)^2 - 6 \cdot 2^x + 8 \leq 0$

i.e.  $(2^x - 4)(2^x - 2) \leq 0$

$2^x \geq 2$  and  $2^x \leq 4$  (Verify !)

$2^x \geq 2^1$  and  $2^x \leq 2^2$

$x \geq 1$  and  $x \leq 2$  or  $1 \leq x \leq 2$

Domain is  $[1,2]$

**8) Logarithmic Function:**

Let,  $a > 0, a \neq 1$ , we define

$y = \log_a x$  if  $x = a^y$ .

for  $x > 0$ , is defined as


$$y = \log_a x \Leftrightarrow a^y = x$$

logarithmic form                      exponential form

**Properties:**

- As  $a^0 = 1$ , so  $\log_a 1 = 0$  and as  $a^1 = a$ , so  $\log_a a = 1$
- As  $a^x = a^y \Leftrightarrow x = y$  so  $\log_a x = \log_a y \Leftrightarrow x = y$
- Product rule of logarithms.  
For  $a, b, c > 0$  and  $a \neq 1$ ,  
 $\log_a bc = \log_a b + \log_a c$  (Verify !)
- Quotient rule of logarithms.  
For  $a, b, c > 0$  and  $a \neq 1$ ,  
 $\log_a \frac{b}{c} = \log_a b - \log_a c$  (Verify !)
- Power/Exponent rule of logarithms.  
For  $a, b, c > 0$  and  $a \neq 1$ ,  
 $\log_a b^c = c \log_a b$  (Verify !)

- 6) For natural base  $e$ ,  $\log_e x = \ln x$  as Natural Logarithm Function.



**Fig. 6.30**

Here domain of  $\ln x$  is  $(0, \infty)$  and range is  $(-\infty, \infty)$ .

- 8) Logarithmic inequalities:

- (i) If  $a > 1$ ,  $0 < m < n$  then  $\log_a m < \log_a n$   
e.g.  $\log_{10} 20 < \log_{10} 30$
- (ii) If  $0 < a < 1$ ,  $0 < m < n$  then  $\log_a m > \log_a n$   
e.g.  $\log_{0.1} 20 > \log_{0.1} 30$
- (iii) For  $a, m > 0$  if  $a$  and  $m$  lies on the same side of unity (i.e. 1) then  $\log_a m > 0$ .  
e.g.  $\log_2 3 > 0$ ,  $\log_{0.3} 0.5 > 0$
- (iv) For  $a, m > 0$  if  $a$  and  $m$  lies on the different sides of unity (i.e. 1) then  $\log_a m < 0$ .  
e.g.  $\log_{0.2} 3 < 0$ ,  $\log_3 0.5 < 0$

**Ex. 11 :** Write  $\log 72$  in terms of  $\log 2$  and  $\log 3$ .

**Solution :**  $\log 72 = \log(2^3 \cdot 3^2)$   
 $= \log 2^3 + \log 3^2$  ( $\because$  Power rule)  
 $= 3 \log 2 + 2 \log 3$  ( $\because$  Power rule)

**Ex. 12 :** Evaluate  $\ln e^9 - \ln e^4$ .

**Solution :**  $\ln e^9 - \ln e^4 = \log_e e^9 - \log_e e^4$   
 $= 9 \log_e e - 4 \log_e e$   
 $= 9(1) - 4(1)$  ( $\because \ln e = 1$ )  
 $= 5$

**Ex. 13 :** Expand  $\log \left[ \frac{x^3(x+3)}{2(x-4)^2} \right]$

**Solution :** Using Quotient rule  
 $= \log [x^3(x+3)] - \log [2(x-4)^2]$   
 Using Product rule  
 $= [\log x^3 + \log(x+3)] - [\log 2 + \log(x-4)^2]$   
 Using Power rule  
 $= [3 \log x + \log(x+3)] - [\log 2 + 2 \log(x-4)]$   
 $= 3 \log x + \log(x+3) - \log 2 + 2 \log(x-4)$

**Ex. 14 :** Combine

$3 \ln(p+1) - \frac{1}{2} \ln r + 5 \ln(2q+3)$  into single logarithm.

**Solution :** Using Power rule,  
 $= \ln(p+1)^3 - \ln r^{\frac{1}{2}} + \ln(2q+3)^5$   
 Using Quotient rule  
 $= \ln \frac{(p+1)^3}{\sqrt{r}} + \ln(2q+3)^5$   
 Using Product rule  
 $= \ln \left[ \frac{(p+1)^3}{\sqrt{r}} (2q+3)^5 \right]$

**Ex. 15 :** Find the domain of  $\ln(x-5)$ .

**Solution :** As  $\ln(x-5)$  is defined for  $(x-5) > 0$  that is  $x > 5$  so domain is  $(5, \infty)$ .

**Let's note:**

- 1)  $\log(x + y) \neq \log x + \log y$
- 2)  $\log x \log y \neq \log(xy)$
- 3)  $\frac{\log x}{\log y} \neq \log\left(\frac{x}{y}\right)$
- 4)  $(\log x)^n \neq n \log x$

9) Change of base formula:

For  $a, x, b > 0$  and  $a, b \neq 1$ ,  $\log_a x = \frac{\log_b x}{\log_b a}$

Note:  $\log_a x = \frac{\log_x x}{\log_x a} = \frac{1}{\log_x a}$  (Verify !)

**Ex. 16 :** Evaluate  $\frac{\log_4 81}{\log_4 9}$

**Solution :** By Change of base law, as the base is same (that is 4)

$$\frac{\log_4 81}{\log_4 9} = \log_9 81 = 2$$

**Ex. 17 :** Prove that,  $2\log_b a^4 \cdot \log_c b^3 \cdot \log_a c^5 = 120$

**Solution :** L.H.S. =  $2\log_b a^4 \cdot \log_c b^3 \cdot \log_a c^5$   
 $= 4 \times 2\log_b a \times 3\log_c b \times 5\log_a c$

Using change of base law,

$$= 4 \times 2 \frac{\log a}{\log b} \times 3 \frac{\log b}{\log c} \times 5 \frac{\log c}{\log a}$$


$$= 120$$

**Ex. 18 :** Find the domain of  $f(x) = \log_{x+5} (x^2 - 4)$

**Solution :** Since  $\log_a x$  is defined for  $a, x > 0$  and  $a \neq 1$   $f(x)$  is defined for  $(x^2 - 4) > 0, x + 5 > 0, x + 5 \neq 1$ .

i.e.  $(x - 2)(x + 2) > 0, x > -5, x \neq -4$

i.e.  $x < -2$  or  $x > 2$  and  $x > -5$  and  $x \neq -4$



**Fig. 6.31**

**9) Trigonometric function**


The graphs of trigonometric functions are discuss in chapter 2 of Mathematics Book I.

$f(x)$	Domain	Range
$\sin x$	R	$[-1, 1]$
$\cos x$	R	$[-1, 1]$
$\tan x$	$R - \left\{ \pm \frac{\pi}{2}, \pm \frac{3\pi}{2} \dots \right\}$	R

**EXERCISE 6.1**


1) Check if the following relations are functions.

(a)




**Fig. 6.32**

(b)



**Fig. 6.33**

(c)



**Fig. 6.34**

2) Which sets of ordered pairs represent functions from  $A = \{1, 2, 3, 4\}$  to  $B = \{-1, 0, 1, 2, 3\}$ ? Justify.

- (a)  $\{(1,0), (3,3), (2,-1), (4,1), (2,2)\}$
- (b)  $\{(1,2), (2,-1), (3,1), (4,3)\}$
- (c)  $\{(1,3), (4,1), (2,2)\}$
- (d)  $\{(1,1), (2,1), (3,1), (4,1)\}$

3) Check if the relation given by the equation represents  $y$  as function of  $x$ .

- (a)  $2x + 3y = 12$
- (b)  $x + y^2 = 9$
- (c)  $x^2 - y = 25$
- (d)  $2y + 10 = 0$
- (e)  $3x - 6 = 21$

4) If  $f(m) = m^2 - 3m + 1$ , find

- (a)  $f(0)$
- (b)  $f(-3)$
- (c)  $f\left(\frac{1}{2}\right)$
- (d)  $f(x+1)$
- (e)  $f(-x)$
- (f)  $\left(\frac{f(2+h) - f(2)}{h}\right), h \neq 0$ .

5) Find  $x$ , if  $g(x) = 0$  where

- (a)  $g(x) = \frac{5x-6}{7}$
- (b)  $g(x) = \frac{18-2x^2}{7}$
- (c)  $g(x) = 6x^2 + x - 2$
- (d)  $g(x) = x^3 - 2x^2 - 5x + 6$

6) Find  $x$ , if  $f(x) = g(x)$  where

- (a)  $f(x) = x^4 + 2x^2, g(x) = 11x^2$
- (b)  $f(x) = \sqrt{x} - 3, g(x) = 5 - x$

7) If  $f(x) = \frac{a-x}{b-x}$ ,  $f(2)$  is undefined, and  $f(3) = 5$ , find  $a$  and  $b$ .

8) Find the domain and range of the following functions.

(a)  $f(x) = 7x^2 + 4x - 1$

(b)  $g(x) = \frac{x+4}{x-2}$

(c)  $h(x) = \frac{\sqrt{x+5}}{5+x}$

(d)  $f(x) = \sqrt[3]{x+1}$

(e)  $f(x) = \sqrt{(x-2)(5-x)}$

(f)  $f(x) = \sqrt{\frac{x-3}{7-x}}$

(g)  $f(x) = \sqrt{16-x^2}$

9) Express the area  $A$  of a square as a function of its (a) side  $s$  (b) perimeter  $P$ .

10) Express the area  $A$  of circle as a function of its (a) radius  $r$  (b) diameter  $d$  (c) circumference  $C$ .

11) An open box is made from a square of cardboard of 30 cms side, by cutting squares of length  $x$  centimeters from each corner and folding the sides up. Express the volume of the box as a function of  $x$ . Also find its domain.

Let  $f$  be a subset of  $Z \times Z$  defined by

12)  $f = \{(ab, a+b) : a, b \in Z\}$ . Is  $f$  a function from  $Z$  to  $Z$ ? Justify.

14) Check the injectivity and surjectivity of the following functions.

(a)  $f: N \rightarrow N$  given by  $f(x) = x^2$

(b)  $f: Z \rightarrow Z$  given by  $f(x) = x^2$

(c)  $f: R \rightarrow R$  given by  $f(x) = x^2$

- (d)  $f: \mathbb{N} \rightarrow \mathbb{N}$  given by  $f(x) = x^3$   
 (e)  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = x^3$
- 14) Show that if  $f: A \rightarrow B$  and  $g: B \rightarrow C$  are one-one, then  $g \circ f$  is also one-one.
- 15) Show that if  $f: A \rightarrow B$  and  $g: B \rightarrow C$  are onto, then  $g \circ f$  is also onto.
- 16) If  $f(x) = 3(4^{x+1})$  find  $f(-3)$ .
- 17) Express the following exponential equations in logarithmic form
- (a)  $2^5 = 32$                       (b)  $54^0 = 1$   
 (c)  $23^1 = 23$                       (d)  $9^{3/2} = 27$   
 (e)  $3^{-4} = \frac{1}{81}$                       (f)  $10^{-2} = 0.01$   
 (g)  $e^2 = 7.3890$                       (h)  $e^{1/2} = 1.6487$   
 (i)  $e^{-x} = 6$
- 18) Express the following logarithmic equations in exponential form
- (a)  $\log_2 64 = 6$                       (b)  $\log_5 \frac{1}{25} = -2$   
 (c)  $\log_{10} 0.001 = -3$                       (d)  $\log_{1/2} (-8) = 3$   
 (e)  $\ln 1 = 0$                       (f)  $\ln e = 1$   
 (g)  $\ln \frac{1}{2} = -0.693$
- 19) Find the domain of
- (a)  $f(x) = \ln(x-5)$   
 (b)  $f(x) = \log_{10}(x^2 - 5x + 6)$
- 20) Write the following expressions as sum or difference of logarithms
- (a)  $\log \left( \frac{pq}{rs} \right)$                       (b)  $\log (\sqrt{x} \sqrt[3]{y})$   
 (c)  $\ln \left( \frac{a^3(a-2)^2}{\sqrt{b^2+5}} \right)$   
 (d)  $\ln \left[ \frac{\sqrt[3]{x-2}(2x+1)^4}{(x+4)\sqrt{2x+4}} \right]^2$
- 21) Write the following expressions as a single logarithm.
- (a)  $5\log x + 7\log y - \log z$   
 (b)  $\frac{1}{3} \log(x-1) + \frac{1}{2} \log(x)$   
 (c)  $\ln(x+2) + \ln(x-2) - 3\ln(x+5)$
- 22) Given that  $\log 2 = a$  and  $\log 3 = b$ , write  $\log \sqrt{96}$  in terms of  $a$  and  $b$ .
- 23) Prove that
- (a)  $b^{\log_b a} = a$                       (b)  $\log_{b^m} a = \frac{1}{m} \log_b a$   
 (c)  $a^{\log_c b} = b^{\log_c a}$
- 24) If  $f(x) = ax^2 - bx + 6$  and  $f(2) = 3$  and  $f(4) = 30$ , find  $a$  and  $b$
- 25) Solve for  $x$ .
- (a)  $\log 2 + \log(x+3) - \log(3x-5) = \log 3$   
 (b)  $2\log_{10} x = 1 + \log_{10} \left( x + \frac{11}{10} \right)$   
 (c)  $\log_2 x + \log_4 x + \log_{16} x = \frac{21}{4}$   
 (d)  $x + \log_{10}(1+2^x) = x \log_{10} 5 + \log_{10} 6$
- 26) If  $\log \left( \frac{x+y}{3} \right) = \frac{1}{2} \log x + \frac{1}{2} \log y$ , show that  $\frac{x}{y} + \frac{y}{x} = 7$ .
- 27) If  $\log \left( \frac{x-y}{4} \right) = \log \sqrt{x} + \log \sqrt{y}$ , show that  $(x+y)^2 = 20xy$
- 28) If  $x = \log_a bc$ ,  $y = \log_b ca$ ,  $z = \log_c ab$  then prove that  $\frac{1}{1+x} + \frac{1}{1+y} + \frac{1}{1+z} = 1$

## 6.2 Algebra of functions:

Let  $f$  and  $g$  be functions with domains  $A$  and  $B$ . Then the functions  $f + g, f - g, fg, \frac{f}{g}$  are defined on  $A \cap B$  as follows.

Operations
$(f + g)(x) = f(x) + g(x)$
$(f - g)(x) = f(x) - g(x)$
$(f \cdot g)(x) = f(x) \cdot g(x)$
$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$ where $g(x) \neq 0$

**Ex. 1 :** If  $f(x) = x^2 + 2$  and  $g(x) = 5x - 8$ , then find

- $(f + g)(1)$
- $(f - g)(-2)$
- $(f \circ g)(3m)$
- $\frac{f}{g}(0)$

**Solution :** i) As  $(f + g)(x) = f(x) + g(x)$

$$\begin{aligned}(f + g)(1) &= f(1) + g(1) \\ &= [(1)^2 + 2] + [5(1) - 8] \\ &= 3 + (-3) \\ &= 0\end{aligned}$$

ii) As  $(f - g)(x) = f(x) - g(x)$

$$\begin{aligned}(f - g)(-2) &= f(-2) - g(-2) \\ &= [(-2)^2 + 2] - [5(-2) - 8] \\ &= [4 + 2] - [-10 - 8] \\ &= 6 + 18 \\ &= 24\end{aligned}$$

iii) As  $(fg)(x) = f(x)g(x)$

$$\begin{aligned}(f \circ g)(3m) &= f(3m)g(3m) \\ &= [(3m)^2 + 2][5(3m) - 8] \\ &= [9m^2 + 2][15m - 8] \\ &= 135m^3 - 72m^2 + 30m - 16\end{aligned}$$

iv) As  $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, g(x) \neq 0$

$$\begin{aligned}\left(\frac{f}{g}\right)(0) &= \frac{f(0)}{g(0)} = \frac{0^2 + 2}{5(0) - 8} \\ &= \frac{2}{-8} = -\frac{1}{4}\end{aligned}$$

**Ex. 2 :** Given the function  $f(x) = 5x^2$  and

$g(x) = \sqrt{4-x}$  find the domain of

- $(f + g)(x)$
- $(f \circ g)(x)$
- $\frac{f}{g}(x)$

**Solution :** i) Domain of  $f(x) = 5x^2$  is  $(-\infty, \infty)$ .

To find domain of  $g(x) = \sqrt{4-x}$

$$\begin{aligned}4 - x &\geq 0 \\ x - 4 &\leq 0\end{aligned}$$

Let  $x \leq 4$ , So domain is  $(-\infty, 4]$ .

Therefore, domain of  $(f + g)(x)$  is  $(-\infty, \infty) \cap (-\infty, 4]$ , that is  $(-\infty, 4]$


ii) Similarly, domain of  $(f \circ g)(x) = 5x^2\sqrt{4-x}$  is  $(-\infty, 4]$

iii) And domain of  $\left(\frac{f}{g}\right)(x) = \frac{5x^2}{\sqrt{4-x}}$  is  $(-\infty, 4)$

As, at  $x = 4$  the denominator  $g(x) = 0$ .

### 6.2.1 Composition of Functions:

A method of combining the function  $f: A \rightarrow B$  with  $g: B \rightarrow C$  is composition of functions, defined as  $(f \circ g)(x) = f[g(x)]$  an read as 'f composed with g'



**Fig. 6.35**

**Note:**

- 1) The domain of  $g \circ f$  is the set of all  $x$  in  $A$  such that  $f(x)$  is in the  $B$ . The range of  $g \circ f$  is set of all  $g[f(x)]$  in  $C$  such that  $f(x)$  is in  $B$ .
- 2) Domain of  $g \circ f \subseteq$  Domain of  $f$  and Range of  $g \circ f \subseteq$  Range of  $g$ .

**Illustration:**

A cow produces 4 liters of milk in a day. Then  $x$  number of cows produce  $4x$  liters of milk in a day. This is given by function  $f(x) = 4x = 'y'$ . Price of one liter milk is Rs. 50. Then the price of  $y$  liters of the milk is Rs.  $50y$ . This is given by another function  $g(y) = 50y$ . Now a function  $h(x)$  gives the money earned from  $x$  number of cows in a day as a composite function of  $f$  and  $g$  as  $h(x) = (g \circ f)(x) = g[f(x)] = g(4x) = 50(4x) = 200x$ .

**Ex. 3 :** If  $f(x) = \frac{2}{x+5}$  and  $g(x) = x^2 - 1$ , then find

- i)  $(f \circ g)(x)$  ii)  $(g \circ f)(3)$

**Solution :**

i) As  $(f \circ g)(x) = f[g(x)]$  and  $f(x) = \frac{2}{x+5}$

Replace  $x$  from  $f(x)$  by  $g(x)$ , to get

$$\begin{aligned} (f \circ g)(x) &= \frac{2}{g(x)+5} \\ &= \frac{2}{x^2-1+5} \\ &= \frac{2}{x^2+4} \end{aligned}$$

ii) As  $(g \circ f)(x) = g[f(x)]$  and  $g(x) = x^2 - 1$   
Replace  $x$  by  $f(x)$ , to get

$$\begin{aligned} (g \circ f)(x) &= [f(x)]^2 - 1 \\ &= \left(\frac{2}{x+5}\right)^2 - 1 \end{aligned}$$

Now let  $x = 3$

$$\begin{aligned} (g \circ f)(3) &= \left(\frac{2}{3+5}\right)^2 - 1 \\ &= \left(\frac{2}{8}\right)^2 - 1 \\ &= \left(\frac{1}{4}\right)^2 - 1 \\ &= \frac{1-16}{16} \\ &= -\frac{15}{16} \end{aligned}$$

**Ex 4 :** If  $f(x) = x^2$ ,  $g(x) = x + 5$ , and  $h(x) = \frac{1}{x}$ ,  $x \neq 0$ , find  $(g \circ f \circ h)(x)$

**Solution :**  $(g \circ f \circ h)(x)$

$$\begin{aligned} &= g\{f[h(x)]\} \\ &= g\left[f\left(\frac{1}{x}\right)\right] \\ &= g\left[\left(\frac{1}{x}\right)^2\right] \\ &= \left(\frac{1}{x}\right)^2 + 5 \\ &= \frac{1}{x^2} + 5 \end{aligned}$$

**Ex. 5 :** If  $h(x) = (x - 5)^2$ , find the functions  $f$  and  $g$ , such that  $h = f \circ g$ .

→ In  $h(x)$ , 5 is subtracted from  $x$  first and then squared. Let  $g(x) = x - 5$  and  $f(x) = x^2$ , (verify)

**Ex. 6 :** Express  $m(x) = \frac{1}{x^3+7}$  in the form of  $f \circ g \circ h$

→ In  $m(x)$ ,  $x$  is cubed first then 7 is added and then its reciprocal taken. So,




$$h(x) = x^3, g(x) = x + 7 \text{ and } f(x) = \frac{1}{x}, \text{ (verify)}$$

### 6.2.2 Inverse functions:

Let  $f : A \rightarrow B$  be one-one and onto function and  $f(x) = y$  for  $x \in A$ . The inverse function

$f^{-1} : B \rightarrow A$  is defined as  $f^{-1}(y) = x$  if  $f(x) = y$



**Fig. 6.36**

#### Note:

- As  $f$  is one-one and onto every element  $y \in B$  has a unique element  $x \in A$  such that  $y = f(x)$ .
- If  $f$  and  $g$  are one-one and onto functions such that  $f[g(x)] = x$  for every  $x \in \text{Domain of } g$  and  $g[f(x)] = x$  for every  $x \in \text{Domain of } f$ , then  $g$  is called inverse of function  $f$ . Function  $g$  is denoted by  $f^{-1}$  (read as  $f$  inverse).  
i.e.  $f[g(x)] = g[f(x)] = x$  then  $g = f^{-1}$  which  
Moreover this means  $f[f^{-1}(x)] = f^{-1}[f(x)] = x$
- $f^{-1}(x) \neq [f(x)]^{-1}$ , because  $[f(x)]^{-1} = \frac{1}{f(x)}$   
 $[f(x)]^{-1}$  is reciprocal of function  $f(x)$  where as  
 $f^{-1}(x)$  is the inverse function of  $f(x)$ .

e.g. If  $f$  is one-one onto function with  $f(3) = 7$  then  $f^{-1}(7) = 3$ .

**Ex. 7 :** If  $f$  is one-one onto function with  $f(x) = 9 - 5x$ , find  $f^{-1}(-1)$ .

**Soln. :**  $\rightarrow$  Let  $f^{-1}(-1) = m$ , then  $-1 = f(m)$

Therefore,

$$-1 = 9 - 5m$$

$$5m = 9 + 1$$

$$5m = 10$$

$$m = 2$$

That is  $f(2) = -1$ , so  $f^{-1}(-1) = 2$ .

**Ex. 8 :** Verify that  $f(x) = \frac{x-5}{8}$  and  $g(x) = 8x + 5$

are inverse functions of each other.

**Solution :** As  $f(x) = \frac{x-5}{8}$ , replace  $x$  in  $f(x)$  with  $g(x)$

$$f[g(x)] = \frac{g(x)-5}{8} = \frac{8x+5-5}{8} = \frac{8x}{8} = x$$

and  $g(x) = 8x + 5$ , replace  $x$  in  $g(x)$  with  $f(x)$

$$g[f(x)] = 8f(x) + 5 = 8 \left[ \frac{x-5}{8} \right] + 5 = x - 5 + 5 = x$$

As  $f[g(x)] = x$  and  $g[f(x)] = x$ ,  $f$  and  $g$  are inverse functions of each other.

**Ex. 9 :** Determine whether the function

$$f(x) = \frac{2x+1}{x-3} \text{ has inverse, if it exists find it.}$$

**Solution :**  $f^{-1}$  exists only if  $f$  is one-one and onto.

$$\text{Consider } f(x_1) = f(x_2),$$

Therefore,

$$\frac{2x_1+1}{x_1-3} = \frac{2x_2+1}{x_2-3}$$

$$(2x_1+1)(x_2-3) = (2x_2+1)(x_1-3)$$

$$2x_1x_2 - 6x_1 + x_2 - 3 = 2x_1x_2 - 6x_2 + x_1 - 3$$

$$-6x_1 + x_2 = -6x_2 + x_1$$

$$6x_1 + x_2 = 6x_2 + x_1$$

$$7x_2 = 7x_1$$

$$x_2 = x_1$$

Hence,  $f$  is one-one function.

$$\text{Let } f(x) = y, \text{ so } y = \frac{2x+1}{x-3}$$

Express  $x$  as function of  $y$ , as follows

$$y = \frac{2x+1}{x-3}$$

$$y(x-3) = 2x+1$$

$$xy - 3y = 2x + 1$$

$$xy - 2x = 3y + 1$$

$$x(y - 2) = 3y + 1$$

$$\therefore x = \frac{3y+1}{y-2} \text{ for } y \neq 2.$$

Thus for any  $y \neq 2$ ,

we have  $x$  such that  $f(x) = y$

$f^{-1}$  is well defined on  $\mathbb{R} - \{2\}$

If  $f(x) = 2$  i.e.  $2x + 1 = 2(x - 3)$

i.e.  $2x + 1 = 2x - 6$  i.e.  $1 = -6$

Which is contradiction.

$2 \notin \text{Range of } f$ .

Here the range of  $f(x)$  is  $\mathbb{R} - \{2\}$ .

$x$  is defined for all  $y$  in the range.

Therefore  $f(x)$  is onto function.

As  $f$  is one-one and onto, so  $f^{-1}$  exists.

As  $f(x) = y$ , so  $f^{-1}(y) = x$

Therefore,  $f^{-1}(y) = \frac{3y+1}{y-2}$

Replace  $x$  by  $y$ , to get

$$f^{-1}(x) = \frac{3x+1}{x-2}.$$

### 6.2.3 Piecewise Defined Functions:

A function defined by two or more equations on different parts of the given domain is called piecewise defined function.

$$\text{e.g.: If } f(x) = \begin{cases} x+1 & \text{if } x < 1 \\ 4-x & \text{if } x \geq 1 \end{cases}$$

Here  $f(3) = 4 - 3 = 1$  as  $3 > 1$ ,

whereas  $f(-2) = -2 + 1 = -1$  as  $-2 < 1$  and

$f(1) = 4 - 1 = 3$ .




Fig. 6.37

As  $(1,3)$  lies on line  $y = 4 - x$ , so it is shown by small black disc on that line.  $(1,2)$  is shown by small white disc on the line  $y = x + 1$ , because it is not on the line.

### 1) Signum function :

**Definition:**  $f(x) = \text{sgn}(x)$  is a piecewise function

$$f(x) = \begin{cases} -1 & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ 1 & \text{if } x > 0 \end{cases}$$




Fig. 6.38

**Domain:**  $\mathbb{R}$  and **Range:**  $\{-1, 0, 1\}$


**Properties:**

- 1) For  $x > 0$ , the graph is line  $y = 1$  and for  $x < 0$ , the graph is line  $y = -1$ .
- 2) For  $f(0) = 0$ , so point  $(0,0)$  is shown by black disc, whereas points  $(0,-1)$  and  $(0,1)$  are shown by white discs.

**Absolute value function (Modulus function):**

Definition:  $f(x) = |x|$ , is a piece wise function

$$f(x) = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$




**Fig. 6.39**

**Domain :**  $\mathbb{R}$  or  $(-\infty, \infty)$  and **Range :**  $[0, \infty)$

**Properties:**

- 1) Graph of  $f(x) = |x|$  is union of line  $y = x$  from quadrant I with the line  $y = -x$  from quadrant II. As origin marks the change of directions of the two lines, we call it a critical point.
- 2) Graph is symmetric about y-axis.
- 3) Graph of  $f(x) = |x-3|$  is the graph of  $|x|$  shifted 3 units right and the critical point is  $(3,0)$ .
- 4)  $f(x) = |x|$ , represents the distance of  $x$  from origin.
- 5) If  $|x| = m$ , then it represents every  $x$  whose distance from origin is  $m$ , that is  $x = +m$  or  $x = -m$ .




**Fig. 6.40**

- 6) If  $|x| < m$ , then it represents every  $x$  whose distance from origin is less than  $m$ ,  $0 \leq x < m$  and  $0 \geq x > -m$  That is  $-m < x < m$ . In interval notation  $x \in (-m, m)$




**Fig. 6.41**

- 7) If  $|x| \geq m$ , then it represents every  $x$  whose distance from origin is greater than or equal to  $m$ , so,  $x \geq m$  and  $x \leq -m$ . In interval notation  $x \in (-\infty, m] \cup [m, \infty)$



**Fig. 6.42**

- 8) If  $m < |x| < n$ , then it represents all  $x$  whose distance from origin is greater than  $m$  but less than  $n$ . That is  $x \in (-n, -m) \cup (m, n)$ .



**Fig. 6.43**

- 9) Triangle inequality  $|x + y| \leq |x| + |y|$ .  
Verify by taking different values for  $x$  and  $y$  (positive or negative).
- 10)  $|x|$  can also be defined as  $|x| = \sqrt{x^2} = \max\{x, -x\}$ .

**Ex. 10 :** Solve  $|4x - 5| \leq 3$ .

**Solution :** If  $|x| \leq m$ , then  $-m \leq x \leq m$

Therefore

$$-3 \leq 4x - 5 \leq 3$$

$$-3 + 5 \leq 4x \leq 3 + 5$$

$$2 \leq 4x \leq 8$$

$$\frac{2}{4} \leq x \leq \frac{8}{4}$$

$$\frac{1}{2} \leq x \leq 2$$

**Ex. 11 :** Find the domain of  $\frac{1}{\sqrt{||x|-1|-3}}$

**Solution :** As function is defined for  $||x|-1|-3 > 0$

Therefore  $||x|-1| > 3$

So  $|x|-1 > 3$  or  $|x|-1 < -3$

That is

$|x| > 3 + 1$  or  $|x| < -3 + 1$

$|x| > 4$  or  $|x| < -2$

But  $|x| < -2$  is not possible as  $|x| > 0$  always


So  $-4 < x < 4$ ,  $x \in (-4, 4)$ .

**Ex. 12 :** Solve  $|x - 1| + |x + 2| = 8$ .

**Solution :** Let  $f(x) = |x - 1| + |x + 2|$

Here the critical points are at  $x = 1$  and  $x = -2$ .

They divide number line into 3 parts, as follows.



**Fig. 6.44**

Region	Test Value	Sign	$f(x)$
I $x < -2$	-3	$(x-1) < 0,$ $(x+2) < 0$	$-(x-1) - (x+2)$ $= -2x - 1$
II $-2 \leq x \leq 1$	0	$(x-1) < 0,$ $(x+2) > 0$	$-(x-1) + (x+2)$ $= 3$
III $x > 1$	2	$(x-1) > 0,$ $(x+2) > 0$	$(x-1) + (x+2)$ $= 2x + 1$

As  $f(x) = 8$

From I,  $-2x - 1 = 8 \therefore -2x = 9 \therefore x = -\frac{9}{2}$ .

From II,  $3 = 8$ , which is impossible, hence there is no solution in this region.

From III,  $2x + 1 = 8 \therefore 2x = 7 \therefore x = \frac{7}{2}$ .

Solutions are  $x = -\frac{9}{2}$  and  $x = \frac{7}{2}$ .

### 3) Greatest Integer Function (Step Function):

**Definition:** For every real  $x$ ,  $f(x) = [x] =$  The greatest integer less than or equal to  $x$ .  $[x]$  is also called as floor function and represented by  $\lfloor x \rfloor$ .

#### Illustrations:

1)  $f(5.7) = [5.7] =$  greatest integer less than or equal to 5.7

Integers less than or equal to 5.7 are 5, 4, 3, 2 of which 5 is the greatest.

2)  $f(-6.3) = [-6.3] =$  greatest integer less than or equal to -6.3.

Integers less than or equal to -6.3 are -10, -9, -8, -7 of which -7 is the greatest.

$\therefore [-6.3] = -7$


3)  $f(2) = [2] =$  greatest integer less than or equal to 2 = 2.

4)  $[\pi] = 3$  5)  $[e] = 2$

The function can be defined piece-wise as follows

$f(x) = n$ , if  $n \leq x < n + 1$  or  $x \in [n, n + 1)$ ,  $n \in \mathbb{I}$

$$f(x) = \begin{cases} -2 & \text{if } -2 \leq x < -1 \text{ or } x \in [-2, -1) \\ -1 & \text{if } -1 \leq x < 0 \text{ or } x \in [-1, 0) \\ 0 & \text{if } 0 \leq x < 1 \text{ or } x \in [0, 1) \\ 1 & \text{if } 1 \leq x < 2 \text{ or } x \in [1, 2) \\ 2 & \text{if } 2 \leq x < 3 \text{ or } x \in [2, 3) \end{cases}$$



Graph of  $f(x) = [x]$

**Fig. 6.45**

Domain =  $\mathbb{R}$  and Range =  $\mathbb{I}$  (Set of integers)

**Properties:**

- 1) If  $x \in [2,3), f(x) = 2$  shown by horizontal line. At exactly  $x = 2, f(2) = 2, 2 \in [2,3)$  hence shown by black disc, whereas  $3 \notin [2,3)$  hence shown by white disc.
- 2) Graph of  $y = [x]$  lies in the region bounded by lines  $y = x$  and  $y = x - 1$ . So  $x - 1 \leq [x] < x$
- 3)  $[x] + [-x] = \begin{cases} 0 & \text{if } x \in I \\ -1 & \text{if } x \notin I \end{cases}$

**Ex.**  $[3.4] + [-3.4] = 3 + (-4) = -1$  where  $3.4 \notin I$   
 $[5] + [-5] = 5 + (-5) = 0$  where  $5 \in I$

4)  $[x+n] = [x] + n$ , where  $n \in I$

**Ex.**  $[4.5 + 7] = [11.5] = 11$  and

$$[4.5] + 7 = 4 + 7 = 11$$

**4) Fractional part function:**


**Definition:** For every real  $x, f(x) = \{x\}$  is defined as  $\{x\} = x - [x]$

**Illustrations:**

$$f(4.8) = \{4.8\} = 4.8 - [4.8] = 4.8 - 4 = 0.8$$

$$f(-7.1) = \{-7.1\} = -7.1 - [-7.1] \\ = -7.1 - (-8) = -7.1 + 8 = 0.9$$

$$f(8) = \{8\} = 8 - [8] = 8 - 8 = 0$$



Graph of  $f(x) = \{x\}$

**Fig. 6.46**

Domain =  $\mathbb{R}$  and Range =  $[0,1)$

**Properties:**

- 1) If  $x \in [0,1), f(x) = \{x\} \in [0,1)$  shown by slant line  $y = x$ . At  $x = 0, f(0) = 0, 0 \in [0,1)$  hence shown by black disc, whereas at  $x = 1, f(1) = 1, 1 \notin [0,1)$  hence shown by white disc.
- 2) Graph of  $y = \{x\}$  lies in the region bounded by  $y = 0$  and  $y = 1$ . So  $0 \leq \{x\} < 1$
- 3)  $\{x\} + \{-x\} = \begin{cases} 0 & \text{if } x \in I \\ 1 & \text{if } x \notin I \end{cases}$

**Ex. 13:**  $\{5.2\} + \{-5.2\} = 0.2 + 0.8 = 1$  where  $5.2 \in I$   
 $\{7\} + \{-7\} = 0 + (0) = 0$  where  $7 \in I$

4)  $\{x \pm n\} = \{x\}$ , where  $n \in I$

**Ex. 14 :**  $\{2.8+5\} = \{7.8\} = 0.8$  and  $\{2.8\} = 0.8$   
 $\{2.8 - 5\} = \{-2.2\} = -2.2 - (-2.2) = -2.2 - (-3) = 0.8$  ( $\because \{x\} = x - [x]$ )

**Ex. 15 :** If  $\{x\}$  and  $[x]$  are the fractional part function and greatest integer function of  $x$  respectively. Solve for  $x$ , if  $\{x + 1\} + 2x = 4[x + 1] - 6$ .

**Solution :**  $\{x + 1\} + 2x = 4[x + 1] - 6$

Since  $\{x + n\} = \{x\}$  and  $[x + n] = [x] + n$ , for  $n \in I$ , also  $x = [x] + \{x\}$

$$\therefore \{x\} + 2(\{x\} + [x]) = 4([x] + 1) - 6$$

$$\therefore \{x\} + 2\{x\} + 2[x] = 4[x] + 4 - 6$$

$$\therefore 3\{x\} = 4[x] - 2[x] - 2$$

$$\therefore 3\{x\} = 2[x] - 2 \quad \dots \text{(I)}$$

Since  $0 \leq \{x\} < 1$

$$\therefore 0 \leq 3\{x\} < 3$$

$$\therefore 0 \leq 2[x] - 2 < 3 \quad (\because \text{from I})$$

$$\therefore 0 + 2 \leq 2[x] < 3 + 2$$

$$\therefore 2 \leq 2[x] < 5$$

$$\therefore \frac{2}{2} \leq [x] < \frac{5}{2}$$

$$\therefore 1 \leq [x] < 2.5$$

But as  $[x]$  takes only integer values

$[x] = 1, 2$  since  $[x] = 1 \Rightarrow 1 \leq x < 2$  and  $[x] = 2 \Rightarrow 2 \leq x < 3$

Therefore  $x \in [1, 3)$

**Note:**

1)

Property	$f(x)$
$f(x+y) = f(x) + f(y)$	$kx$
$f(x+y) = f(x)f(y)$	$a^{kx}$
$f(xy) = f(x)f(y)$	$x^n$
$f(xy) = f(x) + f(y)$	$\log x$

2) If  $n(A) = m$  and  $n(B) = n$  then

(a) number of functions from A and B is  $n^m$  (b) for  $m \leq n$ , number of one-one

functions is  $\frac{n!}{(n-m)!}$

(c) for  $m > n$ , number of one-one functions is 0

(d) for  $m \geq n$ , number of onto functions are  $n^m - {}^n C_1(n-1)^m + {}^n C_2(n-2)^m - {}^n C_3(n-3)^m + \dots + (-1)^{n-1} {}^n C_{n-1}$

(e) for  $m < n$ , number of onto functions are 0.

(f) number of constant functions is  $m$ .

3) Characteristic & Mantissa of Common Logarithm  $\log_{10} x$ :

As  $x = [x] + \{x\}$

$\log_{10} x = [\log_{10} x] + \{\log_{10} x\}$

Where, integral part  $[\log_{10} x]$  is called Characteristic & fractional part  $\{\log_{10} x\}$  is called Mantissa.

**Illustration :** For  $\log_{10} 23$ ,

$\log_{10} 10 < \log_{10} 23 < \log_{10} 100$

$\log_{10} 10 < \log_{10} 23 < \log_{10} 10^2$

$\log_{10} 10 < \log_{10} 23 < 2\log_{10} 10$

$1 < \log_{10} 23 < 2 \quad (\because \log_{10} 10 = 1)$

Then  $[\log_{10} 23] = 1$ , hence Characteristic of  $\log_{10} 23$  is 1.

The characteristic of the logarithm of a number N, with 'm' digits in its integral part is 'm-1'.

**Ex. 16 :** Given that  $\log_{10} 2 = 0.3010$ , find the number of digits in the number  $20^{10}$ .

**Solution :** Let  $x = 20^{10}$ , taking  $\log_{10}$  on either sides, we get

$$\begin{aligned} \log_{10} x &= \log_{10} (20^{10}) = 10\log_{10} 20 \\ &= 10\log_{10} (2 \times 10) = 10\{\log_{10} 2 + \log_{10} 10\} \\ &= 10\{\log_{10} 2 + 1\} = 10\{0.3010 + 1\} \\ &= 10(1.3010) = 13.010 \end{aligned}$$

That is characteristic of  $x$  is 13.

So number of digits in  $x$  is  $13 + 1 = 14$

**EXERCISE 6.2**

1) If  $f(x) = 3x + 5$ ,  $g(x) = 6x - 1$ , then find

- (a)  $(f+g)(x)$                       (b)  $(f-g)(x)$
- (c)  $(fg)(x)$                         (d)  $(f/g)(x)$  and its domain.

2) Let  $f: \{2,4,5\} \rightarrow \{2,3,6\}$  and  $g: \{2,3,6\} \rightarrow \{2,4\}$  be given by  $f = \{(2,3), (4,6), (5,2)\}$  and  $g = \{(2,4), (3,4), (6,2)\}$ . Write down  $g \circ f$

3) If  $f(x) = 2x^2 + 3$ ,  $g(x) = 5x - 2$ , then find

- (a)  $f \circ g$                               (b)  $g \circ f$
- (c)  $f \circ f$                               (d)  $g \circ g$

4) Verify that  $f$  and  $g$  are inverse functions of each other, where

(a)  $f(x) = \frac{x-7}{4}$ ,  $g(x) = 4x + 7$

(b)  $f(x) = x^3 + 4$ ,  $g(x) = \sqrt[3]{x-4}$

(c)  $f(x) = \frac{x+3}{x-2}$ ,  $g(x) = \frac{2x+3}{x-1}$

5) Check if the following functions have an inverse function. If yes, find the inverse function.

(a)  $f(x) = 5x^2$

(b)  $f(x) = 8$

(c)  $f(x) = \frac{6x-7}{3}$

(d)  $f(x) = \sqrt{4x+5}$

(e)  $f(x) = 9x^3 + 8$

(f)  $f(x) = \begin{cases} x+7 & x < 0 \\ 8-x & x \geq 0 \end{cases}$

6) If  $f(x) = \begin{cases} x^2 + 3, & x \leq 2 \\ 5x + 7, & x > 2 \end{cases}$ , then find

(a)  $f(3)$

(b)  $f(2)$

(c)  $f(0)$

7) If  $f(x) = \begin{cases} 4x-2, & x \leq -3 \\ 5, & -3 < x < 3 \\ x^2, & x \geq 3 \end{cases}$ , then find

(a)  $f(-4)$

(b)  $f(-3)$

(c)  $f(1)$

(d)  $f(5)$

8) If  $f(x) = 2|x| + 3x$ , then find

(a)  $f(2)$

(b)  $f(-5)$

9) If  $f(x) = 4[x] - 3$ , where  $[x]$  is greatest integer function of  $x$ , then find

(a)  $f(7.2)$

(b)  $f(0.5)$

(c)  $f\left(-\frac{5}{2}\right)$

(d)  $f(2\pi)$ , where  $\pi = 3.14$

10) If  $f(x) = 2\{x\} + 5x$ , where  $\{x\}$  is fractional part function of  $x$ , then find

(a)  $f(-1)$

(b)  $f\left(\frac{1}{4}\right)$

(c)  $f(-1.2)$

(d)  $f(-6)$

11) Solve the following for  $x$ , where  $|x|$  is modulus function,  $[x]$  is greatest integer function,  $\{x\}$  is a fractional part function.

(a)  $|x+4| \geq 5$

(b)  $|x-4| + |x-2| = 3$

(c)  $x^2 + 7|x| + 12 = 0$

(d)  $|x| \leq 3$

(e)  $2|x| = 5$

(f)  $[x + [x + [x]]] = 9$

(g)  $\{x\} > 4$

(h)  $\{x\} = 0$

(i)  $\{x\} = 0.5$

(j)  $2\{x\} = x + [x]$



### Let's Remember

- If  $f:A \rightarrow B$  is a function and  $f(x) = y$ , where  $x \in A$  and  $y \in B$ , then

**Domain** of  $f$  is  $A =$  Set of Inputs = Set of Pre-images = Set of values of  $x$  for which  $y = f(x)$  is defined = Projection of graph of  $f(x)$  on X-axis.

**Range** of  $f$  is  $f(A) =$  Set of Outputs = Set of Images = Set of values of  $y$  for which  $y = f(x)$  is defined = Projection of graph of  $f(x)$  on Y-axis.

**Co-domain** of  $f$  is  $B$ .

- If  $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$  then  $f$  is **one-one** and for every  $y \in B$ , if there exists  $x \in A$  such that  $f(x) = y$  then  $f$  is **onto**.
- If  $f:A \rightarrow B$ .  $g:B \rightarrow C$  then a function  $g \circ f:A \rightarrow C$  is a **composite function**.
- If  $f:A \rightarrow B$ , then  $f^{-1}:B \rightarrow A$  is **inverse function** of  $f$ .
- If  $f:\mathbb{R} \rightarrow \mathbb{R}$  is a real valued function of real variable, the following table is formed.

Type of $f$	Form of $f$	Domain of $f$	Range of $f$
Constant function	$f(x) = k$	$\mathbb{R}$	$k$
Identity function	$f(x) = x$	$\mathbb{R}$	$\mathbb{R}$
Square function	$f(x) = x^2$	$\mathbb{R}$	$[0, \infty)$ or $\mathbb{R}^+$
Cube function	$f(x) = x^3$	$\mathbb{R}$	$\mathbb{R}$
Linear function	$f(x) = ax + b$	$\mathbb{R}$	$\mathbb{R}$
Quadratic function	$f(x) = ax^2 + bx + c$	$\mathbb{R}$	$\left(\frac{4ac - b^2}{4a}, \infty\right)$
Cubic function	$f(x) = ax^3 + bx^2 + cx + d$	$\mathbb{R}$	$\mathbb{R}$
Square root function	$f(x) = \sqrt{x}$	$[0, \infty)$	$[0, \infty)$ or $\mathbb{R}^+$
Cube root function	$f(x) = \sqrt[3]{x}$	$\mathbb{R}$	$\mathbb{R}$
Rational function	$f(x) = \frac{p(x)}{q(x)}$	$\mathbb{R} - \{x \mid q(x) = 0\}$	depends on $p(x)$ and $q(x)$
Exponential function	$f(x) = a^x, a > 1$	$\mathbb{R}$	$(0, \infty)$
Logarithmic function	$f(x) = \log_a x, a > 1$	$(0, \infty)$ or $\mathbb{R}^+$	$\mathbb{R}$
Absolute function	$f(x) =  x $	$\mathbb{R}$	$[0, \infty)$ or $\mathbb{R}^+$
Signum function	$f(x) = \text{sgn}(x)$	$\mathbb{R}$	$\{-1, 0, 1\}$
Greatest Integer function	$f(x) = [x]$	$\mathbb{R}$	$\mathbb{I}$ (set of integers)
Fractional Part function	$f(x) = \{x\}$	$\mathbb{R}$	$[0, 1)$

### MISCELLANEOUS EXERCISE 6

(I) Select the correct answer from given alternatives.

- If  $\log(5x - 9) - \log(x + 3) = \log 2$  then  $x = \dots\dots\dots$   
 A) 3      B) 5      C) 2      D) 7
- If  $\log_{10}(\log_{10}(\log_{10}x)) = 0$  then  $x =$   
 A) 1000      B)  $10^{10}$   
 C) 10      D) 0
- Find  $x$ , if  $2\log_2 x = 4$   
 A) 4, -4      B) 4  
 C) -4      D) not defined
- The equation  $\log_{x^2} 16 + \log_{2x} 64 = 3$  has,  
 A) one irrational solution  
 B) no prime solution  
 C) two real solutions  
 D) one integral solution
- If  $f(x) = \frac{1}{1-x}$ , then  $f(f(f(x)))$  is  
 A)  $x - 1$       B)  $1 - x$       C)  $x$       D)  $-x$





- 14) If  $f(x) = \frac{x+3}{4x-5}$ ,  $g(x) = \frac{3+5x}{4x-1}$  then show that  $(f \circ g)(x) = x$ .
- 15) Let  $f: \mathbb{R} - \{2\} \rightarrow \mathbb{R}$  be defined by  $f(x) = \frac{x^2-4}{x-2}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $g(x) = x+2$ . Ex whether  $f = g$  or not.
- 16) Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be given by  $f(x) = x+5$  for all  $x \in \mathbb{R}$ . Draw its graph.
- 17) Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be given by  $f(x) = x^3 + 1$  for all  $x \in \mathbb{R}$ . Draw its graph.
- 18) For any base show that  $\log(1+2+3) = \log 1 + \log 2 + \log 3$ .
- 19) Find  $x$ , if  $x = 3^{3 \log_3 2}$
- 20) Show that,  $\log |\sqrt{x^2+1} + x| + \log |\sqrt{x^2+1} - x| = 0$
- 21) Show that,  $\log \frac{a^2}{bc} + \log \frac{b^2}{ca} + \log \frac{c^2}{ab} = 0$
- 22) Simplify,  $\log(\log x^4) - \log(\log x)$ .
- 23) Simplify  $\log_{10} \frac{28}{45} - \log_{10} \frac{35}{324} + \log_{10} \frac{325}{432} - \log_{10} \frac{13}{15}$
- 24) If  $\log \left( \frac{a+b}{2} \right) = \frac{1}{2}(\log a + \log b)$ , then show that  $a=b$
- 25) If  $b^2=ac$ . prove that,  $\log a + \log c = 2 \log b$
- 26) Solve for  $x$ ,  $\log_x(8x-3) - \log_x 4 = 2$
- 27) If  $a^2 + b^2 = 7ab$ , show that,  $\log \left( \frac{a+b}{3} \right) = \frac{1}{2} \log a + \frac{1}{2} \log b$
- 28) If  $\log \left( \frac{x-y}{5} \right) = \frac{1}{2} \log x + \frac{1}{2} \log y$ , show that  $x^2 + y^2 = 27xy$ .
- 29) If  $\log_3 [\log_2(\log_3 x)] = 1$ , show that  $x = 6561$ .
- 30) If  $f(x) = \log(1-x)$ ,  $0 \leq x < 1$  show that  $f \left( \frac{1}{1+x} \right) = f(1-x) - f(-x)$
- 31) Without using log tables, prove that  $\frac{2}{5} < \log_{10} 3 < \frac{1}{2}$
- 32) Show that  $7 \log \left( \frac{15}{16} \right) + 6 \log \left( \frac{8}{3} \right) + 5 \log \left( \frac{2}{5} \right) + \log \left( \frac{32}{25} \right) = \log 3$
- 33) Solve :  $\sqrt{\log_2 x^4 + 4 \log_4 \sqrt{\frac{2}{x}}} = 2$
- 34) Find value of  $\frac{3 + \log_{10} 343}{2 + \frac{1}{2} \log_{10} \left( \frac{49}{4} \right) + \frac{1}{2} \log_{10} \left( \frac{1}{25} \right)}$
- 35) If  $\frac{\log a}{x+y-2z} = \frac{\log b}{y+z-2x} = \frac{\log c}{z+x-2y}$ , show that  $abc = 1$ .
- 36) Show that,  $\log_y x^3 \cdot \log_z y^4 \cdot \log_x z^5 = 60$
- 37) If  $\frac{\log_2 a}{4} = \frac{\log_2 b}{6} = \frac{\log_2 c}{3k}$  and  $a^3 b^2 c = 1$  find the value of  $k$ .
- 38) If  $a^2 = b^3 = c^4 = d^5$ , show that  $\log_a bcd = \frac{47}{30}$ .
- 39) Solve the following for  $x$ , where  $|x|$  is modulus function,  $[x]$  is greatest interger function,  $\{x\}$  is a fractional part function.
- a)  $1 < |x-1| < 4$     c)  $|x^2 - x - 6| = x + 2$   
 c)  $|x^2 - 9| + |x^2 - 4| = 5$   
 d)  $-2 < [x] \leq 7$     e)  $2[2x-5] - 1 = 7$   
 f)  $[x^2] - 5[x] + 6 = 0$   
 g)  $[x-2] + [x+2] + \{x\} = 0$   
 h)  $\left[ \frac{x}{2} \right] + \left[ \frac{x}{3} \right] = \frac{5x}{6}$

40) Find the domain of the following functions.

a)  $f(x) = \frac{x^2 + 4x + 4}{x^2 + x - 6}$

b)  $f(x) = \sqrt{x-3} + \frac{1}{\log(5-x)}$

c)  $f(x) = \sqrt{1 - \sqrt{1 - \sqrt{1 - x^2}}}$

d)  $f(x) = x!$

e)  $f(x) = {}^{5-x}P_{x-1}$

f)  $f(x) = \sqrt{x-x^2} + \sqrt{5-x}$

g)  $f(x) = \sqrt{\log(x^2 - 6x + 6)}$

41) Find the range of the following functions.

a)  $f(x) = |x-5|$       b)  $f(x) = \frac{x}{9+x^2}$

c)  $f(x) = \frac{1}{1+\sqrt{x}}$       d)  $f(x) = [x] - x$

e)  $f(x) = 1 + 2^x + 4^x$

42) Find  $(f \circ g)(x)$  and  $(g \circ f)(x)$

a)  $f(x) = e^x, g(x) = \log x$

b)  $f(x) = \frac{x}{x+1}, g(x) = \frac{x}{1-x}$

43) Find  $f(x)$  if

a)  $g(x) = x^2 + x - 2$  and  $(g \circ f)(x) = 4x^2 - 10x + 4$

(b)  $g(x) = 1 + \sqrt{x}$  and  $f[g(x)] = 3 + 2\sqrt{x} + x$ .

44) Find  $(f \circ f)(x)$  if

(a)  $f(x) = \frac{x}{\sqrt{1+x^2}}$

(b)  $f(x) = \frac{2x+1}{3x-2}$

